SCIENXT

ISSN: 2584-1874; SJBLS
Scienxt Journal of Biotechnology and Life Science

Anatomical and physiological perspectives in paediatric homeopathy: a comprehensive review of organ-system development, individualised therapeutics, and clinical applications

Dr. R Imran Khan[1]*, Dr. Sanchita Singh[2]

*[1]*BHMS MD(HOM), Department of Human Anatomy, RRHMC & H, Nelamangala, Bengaluru, India*

[2]Assistant Professor, Department of Human Anatomy, Rosy Royal Homoeopathic Medical College, Bangalore, India

DOI : 10.5281/zenodo.23099878

Abstract:

Paediatric care requires an understanding that anatomy and physiology are dynamic across infancy, childhood and adolescence. Organ size, tissue composition, neural maturation, cardiopulmonary reserve, renal and hepatic function, endocrine transitions, immune development, and behavioural communication all change with age. These differences influence disease presentation, diagnostic thresholds, medicine disposition, treatment response, and safety. Homeopathy is a longstanding complementary medical system in which treatment is individualised around a symptom pattern and traditionally involves highly diluted preparations. The central purpose of this review is not to assume that homeopathy modifies normal organ maturation, but to examine how paediatric developmental science can inform the clinical appraisal of homeopathic practice, the interpretation of symptom patterns, and the design of safe integrative care. Current evidence from systematic reviews and government evidence evaluations is discussed condition by condition. Studies of childhood respiratory infections, atopic disease, neurodevelopmental disorders, gastrointestinal complaints and other conditions remain heterogeneous, and higher-quality evidence has generally not demonstrated a consistent specific effect beyond placebo. The 2024 Australian evidence evaluation also identified substantial uncertainty across many outcomes and emphasised the limitations of the available evidence. Safety deserves equal attention because children have age-dependent physiological vulnerability, limited ability to describe symptoms, and a greater risk of harm when effective conventional treatment is delayed. A developmentally informed approach therefore places growth monitoring, diagnostic assessment, vaccination, nutrition, emergency recognition, evidence-based treatment and communication with caregivers before any complementary intervention. Homeopathy may be considered by some families as an adjunctive approach, but it should not replace proven care for serious or progressive paediatric disease. Future research should combine rigorous randomisation, transparent individualisation protocols, age-stratified outcomes, validated developmental measures, pharmacovigilance and reporting of treatment displacement.

Keywords:

Paediatric homeopathy; child development; anatomy; physiology; individualised therapeutics; complementary medicine; developmental pharmacology; organ-system maturation; clinical safety; evidence-based paediatrics.

1. Introduction:

Childhood is not simply a smaller version of adulthood. The body is continuously changing in structure, composition, function and regulatory capacity. Growth in length and weight is accompanied by changes in body water, extracellular fluid, protein binding, fat mass, organ blood flow, enzyme activity, receptor expression, immune competence and neurocognitive function. These processes occur at different rates, so a child of a given chronological age can differ substantially from another child in biological maturation. The clinical consequence is that the same disease may produce different signs in a neonate, infant, school-aged child and adolescent, and an intervention that is acceptable in an adult may not be appropriate in a younger patient. Developmental pharmacology describes these changes as important determinants of drug disposition and drug response (Kearns et al., 2003; van den Anker et al., 2018).

Homeopathy occupies a distinctive position within complementary medicine because its therapeutic reasoning is based on individualisation, symptom similarity and potentised preparations rather than on conventional pharmacological dose-response relationships. In homeopathic consultations, the clinician may collect a broad narrative that includes presenting symptoms, timing, triggers, food preferences, sleep, emotional context and constitutional features. Such attention to the whole child can overlap with a good paediatric history, but overlap in consultation style does not establish that homeopathic remedies have specific biological efficacy. Modern evidence appraisal separates the quality of the consultation, the natural course of illness, contextual effects, and the specific effect of the administered preparation. This distinction is essential when discussing homeopathy in children because parental expectations and the therapeutic relationship can influence observed outcomes even when a treatment does not outperform placebo (Shang et al., 2005; NCCIH, 2025).

Interest in paediatric homeopathy is partly driven by the high frequency of self-limiting childhood symptoms, concern about medicine exposure, desire for non-pharmacological care, and interest in individualised treatment. However, the same features that make children attractive candidates for “gentle” approaches also create a safety obligation. Infants cannot describe symptoms reliably; deterioration may occur quickly; dehydration can develop rapidly; respiratory reserve is limited in severe illness; and developmental disorders can be missed when symptoms are interpreted only as personality or temperament. Current complementary-medicine guidance therefore recommends that children receive an accurate diagnosis and that complementary interventions should not delay or replace treatments with established safety and effectiveness (NCCIH, 2025a).

Anatomical and physiological perspectives are especially useful because they provide a common clinical language between paediatricians, nurses, pharmacists, allied professionals and complementary practitioners. The anatomy of a small airway, immature renal clearance, changing thermoregulation, evolving immune responses, and the continuing maturation of the brain all influence what constitutes normal and abnormal. The purpose of this article is to integrate these principles with the literature on homeopathy, identify potential areas of clinical application, and define boundaries for responsible use.

2. Aim and Review Approach:

The aim of this review is to provide an integrated, academically oriented overview of paediatric organ-system development and to examine how developmental anatomy and physiology should shape assessment, individualised consultation, clinical application and safety considerations in paediatric homeopathy. The review also examines the current evidence for common paediatric indications and identifies methodological priorities for future research.

A narrative integrative approach was used. Targeted literature was identified from PubMed-indexed biomedical studies, World Health Organization publications, the US National Center for Complementary and Integrative Health, and government evidence evaluations. Priority was given to systematic reviews, meta-analyses, major developmental physiology reviews, official child-growth resources and recent government evidence assessments. The evidence was interpreted conservatively because homeopathy-specific studies often differ in remedy selection, consultation structure, comparator, outcome definition and risk of bias. The article therefore uses phrases such as “has been

studied", "has been reported", and "evidence remains uncertain" when a causal conclusion is not supported.

The review focuses on human paediatric development from infancy through adolescence. "Paediatric homeopathy" is used descriptively to refer to homeopathic care delivered to children and adolescents; it does not imply that homeopathy is a component of established conventional paediatric treatment. The article does not provide individual prescribing instructions, dosing schedules, dilution recommendations or advice to discontinue prescribed medicines. Instead, it emphasises clinical assessment, evidence interpretation, shared decision-making and safeguarding.

3.Paediatric Development as the Anatomical and Physiological Foundation:

3.1. Age, growth and developmental windows:

Paediatric development can be described using overlapping age groups such as newborn, infant, toddler, preschool child, school-aged child and adolescent. These labels are clinically useful but should not be treated as rigid biological boundaries. Weight, length or height, head circumference, body mass index and growth velocity are interpreted against age- and sex-specific references. The WHO 2006 standards for children from birth to 5 years were developed from a multicentre study of approximately 8,400 children and are intended to describe how children grow under optimal conditions (World Health Organization, 2006). For older children and adolescents, WHO recommends growth references from 5 to 19 years (World Health Organization, 2025).

Growth surveillance matters in any setting where parents report poor appetite, recurrent illness, chronic fatigue, altered bowel function, sleep difficulties or developmental concerns. A falling weight-for-age trajectory may indicate inadequate intake, malabsorption, chronic inflammation, endocrine disease or other pathology. Conversely, rapid changes in weight can reflect excess energy intake, endocrine disorders, medication effects or fluid retention. A complementary consultation that records symptoms but does not measure growth can therefore miss a clinically important signal. In paediatric practice, trend is usually more informative than a single number, particularly when growth is assessed alongside height, body mass index, head circumference in younger children, developmental history and pubertal stage.

Development also occurs in time-sensitive windows. Neural connections are refined through synaptic pruning and myelination, lung structure and function continue to mature after birth, and immune exposure helps shape adaptive responses. Adolescence introduces another period of rapid biological change associated with hormonal transitions and continuing neural reorganisation (Taki and Kawashima, 2012; Dow-Edwards et al., 2019). The practical implication for individualised history taking is that age, developmental stage and context should be recorded before symptom meaning is inferred.

4.Development of Individual Organ Systems:

4.1. Nervous system: structure, function and behaviour:

The nervous system undergoes prolonged structural and functional maturation. Brain development is non-linear, with different cortical regions reaching developmental peaks at different times; white-matter volume generally increases across childhood and adolescence, while functional networks continue to refine (Taki and Kawashima, 2012). During adolescence, synaptic pruning, myelination, prefrontal maturation and changes in the integration of limbic and executive networks contribute to changing attention, impulse control and emotional regulation (Dow-Edwards et al., 2019). These processes are normal biology rather than signs of disease.

For paediatric assessment, nervous-system development means that symptoms should be interpreted against age. A toddler with a limited vocabulary cannot be evaluated using the same symptom language as a teenager who can describe mood, intrusive thoughts or headaches. Behaviour is also multidimensional. Sleep deprivation, pain, hearing impairment, school stress, learning difficulties, neurodevelopmental conditions and family circumstances can all influence attention or irritability. A broad homeopathic consultation can capture these domains, but a comprehensive history does not

replace developmental screening or formal assessment when symptoms are persistent, impairing or associated with regression.

In the context of homeopathy, the temptation to treat behavioural characteristics as constitutional markers requires caution. Traits such as restlessness, shyness, perfectionism or sensitivity may be developmentally normal or may occur in several different disorders. Diagnostic labels should therefore be established using accepted clinical criteria before complementary treatment is considered. Altunç et al. found that randomised trials in children and adolescents covered several conditions but concluded that the evidence was not convincing enough to recommend homeopathy for any paediatric condition (Altunç et al., 2007). This remains relevant when homeopathic case narratives are used in neurodevelopmental complaints.

4.2. Cardiovascular system:

The cardiovascular system changes markedly after birth. Transition from placental circulation to independent pulmonary gas exchange alters pulmonary vascular resistance, cardiac loading and the pattern of blood flow. Heart rate is higher in infants and younger children than in adults, and normal values vary with age, sleep, activity and fever. Stroke volume, vascular compliance and autonomic regulation also mature over time. These differences affect how paediatric illness presents and how vital signs should be interpreted.

A homeopathic consultation involving pallor, exercise intolerance, cyanosis, syncope, chest pain, tachycardia or unexplained breathlessness must first distinguish benign variation from potentially significant cardiovascular disease. Congenital heart disease, myocarditis, arrhythmia, anaemia and respiratory disorders can overlap clinically. A child with a new murmur, persistent cyanosis, exertional syncope or signs of poor perfusion requires medical evaluation rather than a trial of complementary treatment. The principle is simple: individualised symptom description is useful only after red flags have been recognised and the underlying physiology has been assessed.

4.3. Respiratory system:

Respiratory development begins during fetal life and continues into early childhood. The airway is smaller in young children, the chest wall is more compliant, respiratory muscles are less developed and the functional reserve available during severe respiratory disease is limited. Lung volumes and mechanics change rapidly during infancy, while alveolar and airway development continue after birth (Stocks, 1977; Mullassery and Smith, 2015). Early adverse influences on lung development can be associated with later respiratory morbidity and reduced lung function (Stocks et al., 2013). These developmental features explain why a seemingly modest increase in airway obstruction can produce significant work of breathing in an infant. Paediatric respiratory assessment therefore prioritises respiratory rate, retractions, nasal flaring, grunting, wheeze, stridor, oxygen saturation where available, hydration and feeding ability. Homeopathic symptom narratives may document the timing of cough, aggravating factors and associated symptoms, but these details do not establish a specific effect of a remedy on bronchial smooth muscle, airway inflammation or gas exchange. Hormonal factors may also influence paediatric pulmonary development and function. Recent review literature highlights complex interactions among endocrine signals, airway tone, inflammatory regulation and sex-related differences across childhood and adolescence (Pelizzo et al., 2024). This reinforces the need to consider developmental stage rather than treating respiratory symptoms as static. In suspected asthma, recurrent wheezing or chronic cough, diagnosis and guideline-based management remain central. The 2022 Cochrane review and the 2021 systematic review of homeopathic treatment and prevention of acute respiratory tract infections in children did not find convincing evidence of benefit in higher-quality trials (Hawke et al., 2022; King et al., 2021).

6. Gastrointestinal system and the developing gut environment:

The gastrointestinal system changes in relation to feeding, motility, enzyme activity, barrier function and microbial colonisation. Infancy is a particularly dynamic period because the diet changes from milk-based feeding to complementary foods, altering substrate availability for the gut microbiota. The developing microbiome is associated with maturation of gastrointestinal, immune and metabolic functions and may communicate with distant organs including the brain and lungs (Li et al., 2014; Ahearn-Ford et al., 2022). From a homeopathic perspective, gastrointestinal symptoms are often

described in great detail because stool character, appetite, food preferences, abdominal distension and symptom timing may be prominent in case taking. Clinically, however, stool description must be integrated with hydration, growth, abdominal examination and red flags. Blood in stool, bilious vomiting, persistent severe abdominal pain, dehydration, weight loss, abdominal distension or recurrent nocturnal symptoms require diagnostic assessment. Homeopathic consultation should never be used to normalise these findings without investigation.

The developing microbiome is scientifically interesting, but the existence of microbiome-host interactions does not establish that homeopathic dilutions specifically modify microbial ecology. Claims about "gut healing", "immune rebalancing" or microbiome correction should therefore be presented as hypotheses unless supported by direct controlled evidence. The strongest current evidence concerns the biology of microbiome development itself, not a proven homeopathic mechanism.

7. Renal system and fluid-electrolyte regulation:

Renal function develops after birth, and the capacity for glomerular filtration, tubular handling and urinary concentration changes with maturation. Young children have a different total body-water distribution and a higher proportion of extracellular water than older children and adults. During vomiting, diarrhoea, fever or poor intake, fluid balance can therefore change rapidly. Renal maturation also has major implications for the use of conventional medicines because clearance may be markedly age-dependent (Kearns et al., 2003; van den Anker et al., 2018). In complementary consultations, urine frequency, colour, thirst and enuresis may be reported as individualising symptoms. These details can support history taking but should not substitute for urinalysis or clinical examination when indicated. Persistent oedema, hypertension, haematuria, significant reduction in urine output or recurrent urinary infection requires conventional evaluation. An anatomically and physiologically informed approach treats the symptom as a clinical sign first and as a potential individualising feature second.

8. Liver, metabolism and developmental pharmacology:

The liver is not fully equivalent to an adult organ at birth. Drug-metabolising enzymes, transporters, plasma proteins and hepatic blood flow change during development. Together with changes in kidney function, gastrointestinal absorption, body composition and receptor systems, these factors influence the exposure and effect of medicines. Developmental pharmacology therefore explains why adult doses cannot simply be scaled linearly by body weight and why paediatric formulations and age-specific dosing studies are needed (Kearns et al., 2003; Russel et al., 2020).

Although highly diluted homeopathic products may contain little or no measurable quantity of the starting material, the category "homeopathic" is not synonymous with "risk-free". Products may differ in dilution, manufacturing quality and ingredients, and some products labelled homeopathic can contain pharmacologically active substances. NCCIH notes that adverse effects and interactions are possible with some homeopathic products (NCCIH, 2025b). This is particularly relevant for infants, children with complex disease, and families using several complementary products simultaneously. The developmental pharmacology literature also provides a useful conceptual lesson for homeopathic research: intervention effect cannot be separated from the biological context of the recipient. Any future clinical study should therefore report age, weight, developmental stage, comorbidity, conventional medicines, nutrition and relevant organ function. Without these variables, apparent treatment differences may be confounded by normal maturation or unequal baseline risk (van den Anker et al., 2018; Chapron et al., 2022).

8.1. Immune and lymphoid development:

Immune development begins before birth and continues throughout early childhood. Exposure to maternal factors, environmental microorganisms, vaccines, nutrition and the gut microbiome influences immune maturation. Reviews describe childhood immune development as an active process in which innate and adaptive responses gradually acquire greater functional complexity (Djuardi et al., 2011). The lymphoid tissues also change with age and exposure history.

The immune system can be misunderstood when recurrent childhood infections are interpreted as evidence of a chronically "weak immune system" without checking exposure, age, childcare attendance, vaccination status, nutrition, anatomy and the severity of infections. Many young children have frequent viral infections because of normal exposure. Recurrent severe, unusual or opportunistic infections, poor growth or persistent fever warrant medical investigation. Homeopathy should not be promoted as a substitute for vaccination or for evaluation of suspected immunodeficiency. This safety boundary is especially important because delayed diagnosis can have consequences beyond symptom duration.

8.2. Endocrine system and puberty:

Endocrine signalling coordinates growth, metabolism, stress responses and reproductive maturation. During childhood, growth hormone and insulin-like growth factor pathways contribute to linear growth, while thyroid hormones support metabolic and neurodevelopmental processes. Puberty introduces activation of the hypothalamic-pituitary-gonadal axis with rapid changes in growth velocity, body composition, secondary sexual characteristics and emotional experience. Because endocrine maturation varies between individuals, assessment should consider family pattern, growth trajectory and pubertal staging rather than relying on a single chronological expectation. Symptoms sometimes presented in homeopathic consultations as constitutional or emotional patterns can overlap with endocrine complaints. Fatigue may occur with sleep deprivation, iron deficiency, thyroid disease, chronic inflammation or psychosocial stress. Weight change can reflect nutrition, endocrine disease or medication exposure. Menstrual irregularity during early adolescence may be part of normal maturation but can also signal endocrine or gynaecological disease. A developmentally informed consultation therefore places the symptom within the appropriate physiological window before assigning therapeutic significance.

8.3. Musculoskeletal system and connective tissues:

Bone, muscle and connective tissues change in size, strength and composition throughout childhood. Growth plates remain active until skeletal maturity, making paediatric injuries and overuse disorders different from those seen in adults. Muscle mass increases gradually, and neuromuscular coordination continues to mature. Nutritional factors such as protein, calcium and vitamin D availability influence growth, while physical activity supports bone and metabolic health.

When children present with recurrent limb pain, joint swelling, gait disturbance or reduced activity, careful assessment is needed to distinguish benign developmental pain from inflammatory, infectious, neoplastic or mechanical disorders. Persistent swelling, morning stiffness, fever, weight loss, night pain or functional decline requires medical investigation. Individualised symptom descriptions may be recorded within a broader consultation, but they should not become a reason to delay diagnostic imaging, laboratory tests or specialist referral when red flags are present.

8.4. Skin, thermoregulation and sensory systems:

Children have a relatively large body surface area compared with adults, and infants have immature thermoregulatory mechanisms. Skin barrier function also develops after birth, while atopic dermatitis reflects a complex interaction among barrier integrity, immune regulation and environmental triggers. Sensory development continues through childhood, and visual and auditory function strongly influence language, learning and behaviour.

Skin symptoms such as eczema, urticaria, recurrent rash or itching are common reasons families seek complementary care. The phenotype should be described carefully, including distribution, itch severity, sleep disturbance, infection signs and treatment history. The controlled evidence base for homeopathy in eczema is particularly weak; a systematic review identified only a small number of controlled trials, all methodologically limited, and found no demonstrated efficacy (Ernst, 2012). Therefore, supportive skin care and evidence-based management should not be displaced by unproven treatment.

9. Foundations of Homeopathy and the Meaning of Individualisation:

Homeopathy developed in Europe more than two centuries ago and is commonly described using two central concepts: "like cures like" and treatment using highly diluted preparations. In many

homeopathic traditions, a consultation seeks to identify an individual symptom pattern and then select a preparation according to a materia medica and repertory framework. NCCIH notes that some products are diluted to the point at which no molecules of the original substance may remain, while also warning that products labelled homeopathic can vary and may contain active ingredients (NCCIH, 2025b).

Individualisation is an important conceptual feature because it changes what the intervention means in research. In conventional drug trials, the active intervention can often be specified before randomisation and given using a reproducible dose. In individualised homeopathy, the consultation can determine which remedy is selected, making the intervention a package of assessment, clinician interaction, remedy choice and follow-up. The 2014 systematic review by Mathie et al. evaluated randomised placebo-controlled trials of individualised homeopathy and reported that the evidence base was heterogeneous and sensitive to methodological considerations (Mathie et al., 2014).

Individualisation should therefore be separated into two questions. The first is whether a comprehensive, person-centred consultation is clinically valuable. The second is whether the selected homeopathic preparation has a specific effect beyond placebo and the contextual effects of the consultation. The first question can often be answered positively as a matter of communication and clinical assessment. The second requires controlled evidence and cannot be inferred from the quality of the consultation itself. For children, individualisation has another dimension: development changes the meaning of symptoms. A sleep pattern, stool frequency, appetite preference or emotional response can be normal at one age and concerning at another. The homeopathic clinician must therefore be able to interpret the same symptom against paediatric norms. In this sense, anatomical and physiological knowledge is not an accessory to paediatric homeopathy; it is necessary to avoid overinterpreting ordinary developmental variation.

10.Individualised Paediatric Assessment: A Developmentally Informed Clinical Framework:

A safe paediatric complementary consultation begins with the same core questions used in responsible clinical practice: What is the child's age and developmental stage? What is the main concern? When did it begin? How severe is it? What is its trajectory? What factors worsen or relieve it? What has already been evaluated or treated? The history should also cover growth, nutrition, sleep, elimination, immunisation, allergies, previous admissions, medications, family history and relevant exposures. For adolescents, confidentiality and autonomy should be handled in accordance with local law and professional standards.

Observation is especially important in younger children. General appearance, hydration, interaction, respiratory effort, skin colour, activity level and feeding can provide information before the child can answer questions. Developmental surveillance should include communication, motor skills, social interaction and school functioning as appropriate. A complementary practitioner who recognises an abnormal finding should document it and facilitate medical assessment instead of interpreting it solely through a homeopathic framework. The next step is risk stratification. Symptoms such as severe breathing difficulty, altered consciousness, persistent cyanosis, seizure, severe dehydration, anaphylaxis, suspected meningitis, significant trauma or suspected poisoning require urgent conventional medical care. A less acute presentation may permit discussion of complementary approaches, but the decision should still consider diagnosis, evidence, potential interactions and the consequences of treatment delay. In homeopathic case analysis, individualising symptoms may be organised into characteristic features, modalities, associated symptoms and broader constitutional history. A clinically responsible framework can retain these details while preventing premature causation. For example, "cough worse at night" is a descriptive symptom; it is not evidence that a specific homeopathic remedy altered bronchial physiology. Likewise, "child becomes restless when fever rises" can be documented without assuming that it represents a constitutional disorder.

Follow-up should use predefined outcomes rather than general impressions alone. Useful measures include symptom frequency, severity scales, sleep duration, school attendance, rescue medicine use when relevant, growth measures, validated quality-of-life instruments, and caregiver-reported functioning. Objective outcomes reduce the risk that improvement is attributed automatically to the

remedy when the disease is naturally resolving. They also help distinguish symptomatic improvement from a change in the underlying disease process.




Figure.1: Developmentally informed pathway for an individualised paediatric homeopathic consultation

11. Clinical Applications: What Has Been Studied in Paediatric Homeopathy?:

The clinical literature on paediatric homeopathy spans acute respiratory infections, recurrent otitis media, allergic and atopic conditions, asthma, gastrointestinal disorders, behavioural or neurodevelopmental complaints, sleep problems, headache and other chronic symptoms. The presence of research does not mean that an indication is established. Evidence must be considered according to study design, sample size, blinding, comparator, outcome validity, risk of bias and consistency across trials.

12.Acute upper respiratory tract infections:

Upper respiratory tract infections are among the most common childhood illnesses and are usually self-limiting. This makes them an important context for evaluating complementary treatments because spontaneous improvement is common, symptom severity varies, and caregiver outcomes such as comfort and sleep may improve over several days regardless of intervention. In their 2021 systematic review and meta-analysis of eight double-blind randomised studies involving 1,562 children, King et al. found that studies at low risk of bias did not show benefit from homeopathy and concluded that effectiveness was not supported in higher-quality trials (King et al., 2021). The 2022 Cochrane review similarly examined homeopathic medicinal products for preventing and treating acute respiratory tract infections in children. Its findings reinforce the need to distinguish a plausible-seeming improvement in a self-limiting illness from a treatment effect that can be replicated under controlled conditions (Hawke et al., 2022). A developmentally informed clinician should also remember that infants with respiratory symptoms may deteriorate faster than older children because of smaller airways and limited reserve.

13. Otitis media and recurrent infections:

Childhood otitis media is strongly influenced by age, Eustachian-tube anatomy, exposure to respiratory pathogens, smoke exposure and childcare settings. Because recurrence is common, parents may perceive fewer episodes after a particular intervention even when recurrence naturally falls with maturation. A recent government evidence evaluation reviewed the homeopathy literature on recurrent childhood otitis media as a distinct evidence category (Maurin et al., 2024). The practical lesson is that an apparent reduction in episodes should be interpreted with attention to the natural

history of the condition, objective diagnostic criteria and the number of episodes before and after treatment. In a child with severe otalgia, persistent fever, mastoid swelling, significant hearing loss, balance disturbance or toxic appearance, the priority is conventional assessment. Hearing and speech development should also be considered because recurrent middle-ear disease can affect communication during critical developmental windows. An individualised consultation may complement communication with families, but it should not replace otoscopic examination or hearing evaluation when indicated.

14. Asthma, recurrent wheeze and allergic rhinitis:

Asthma is a chronic inflammatory airway disorder with variable symptoms and airway obstruction. The paediatric phenotype changes with age, and recurrent wheeze in early childhood is not always equivalent to established asthma. Treatment decisions depend on severity, exacerbation history, triggers, objective assessment where feasible, and guideline-directed use of inhaled medicines. A complementary treatment should not be positioned as a replacement for inhaled corticosteroids, rescue therapy or emergency treatment when these are clinically indicated.

Homeopathy has been studied in asthma and respiratory allergy, but evidence remains uncertain. The 2024 Australian evidence evaluation included asthma among its prespecified clinical categories, while earlier paediatric reviews did not establish convincing benefit (Maurin et al., 2024; Altunç et al., 2007). For allergic rhinitis, Banerjee et al. reviewed controlled clinical evidence and found the literature difficult to interpret because of study limitations and heterogeneity (Banerjee et al., 2017). From an organ-system perspective, a respiratory history can be made more precise by documenting age at first wheeze, interval symptoms, nocturnal symptoms, exercise limitation, feeding effects in infants, exposure to smoke or allergens, and previous hospital treatment. These factors are more clinically informative than a broad claim that the child has an “allergic constitution”.

15.Atopic dermatitis and other skin conditions:

Atopic dermatitis often fluctuates over time, and its clinical course is influenced by skin barrier function, immune processes, irritants, infection, climate, bathing practices, stress and adherence to treatment. Improvement can occur spontaneously or with routine skin care, creating challenges for uncontrolled observational studies. Ernst’s systematic review of controlled trials concluded that the evidence did not demonstrate efficacy of homeopathy for eczema (Ernst, 2012). A responsible complementary consultation can still add value through detailed assessment of triggers, sleep disruption, family concerns, treatment burden and adherence. However, skin infections, extensive involvement, severe pruritus, excoriation, poor sleep or growth effects may require escalation. The aim should be to reduce distress without creating the false impression that homeopathic treatment has replaced topical anti-inflammatory therapy, emollient care, infection management or allergy assessment when appropriate.

16.Gastrointestinal complaints: diarrhoea, colic and functional symptoms:

Acute diarrhoea illustrates the importance of physiology in paediatric complementary care. The immediate clinical problem is not only the number of stools but also fluid and electrolyte losses, reduced intake, vomiting and the child’s ability to maintain hydration. Infants are particularly vulnerable to dehydration. Homeopathic treatment should therefore never substitute for oral rehydration, nutritional support and medical assessment when clinically indicated. The historical paediatric literature on homeopathy for diarrhoea contains mixed findings, which was one reason broader reviews concluded that evidence was not sufficiently convincing (Altunç et al., 2007).

Infantile colic is another area in which parental expectations, regression to the mean and spontaneous resolution can strongly influence perceived treatment benefit. The 2024 Australian evidence evaluation identified infantile colic and other digestive disorders as a separate evidence category (Maurin et al., 2024). Future trials should use validated crying-duration measures, blinded assessment where feasible, prespecified minimum clinically important differences and careful reporting of feeding and growth. For recurrent abdominal pain or constipation, clinicians should consider diet,

stool withholding, psychosocial factors, medications, coeliac disease, inflammatory disease and other diagnoses according to the clinical picture. Homeopathic symptom descriptors can be recorded as part of the narrative, but the diagnosis should be established using recognised paediatric criteria and appropriate examination.

17.Neurodevelopmental, behavioural and sleep-related complaints:

Children and adolescents may present with inattention, hyperactivity, anxiety, sleep disturbance, headaches, school difficulty or emotional dysregulation. These complaints are strongly influenced by brain development, sleep, learning environment, family stress, sensory impairment and comorbidity. A symptom-based complementary framework should therefore avoid reducing complex developmental phenomena to a single remedy picture.

The paediatric evidence for homeopathy in neurodevelopmental disorders is mixed. The review by Altunç et al. identified both positive and negative findings in ADHD trials and concluded that the overall evidence was not convincing enough for recommendations (Altunç et al., 2007). A broader 2024 government evidence evaluation also examined neurodevelopmental disorders and reported substantial uncertainty across the evidence base (Maurin et al., 2024). The presence of a positive individual study should therefore not be interpreted as proof of effectiveness. Sleep complaints deserve separate attention because sleep duration and architecture change with age. When a child is tired, irritable or inattentive, clinicians should assess bedtime routine, snoring, obstructive symptoms, restless sleep, screen exposure, anxiety, school demands and medication effects. A homeopathic consultation may document these features in depth, but persistent sleep problems should be assessed using age-appropriate clinical pathways.

18.Headache, anxiety and other chronic symptoms:

Headache in children can reflect migraine, tension-type headache, infection, refractive problems, sleep disruption, dehydration or, less commonly, neurological disease. A new severe headache, headache with neurological deficit, altered consciousness, persistent vomiting or other red flags requires medical assessment. Chronic symptoms should be tracked with frequency and functional impact rather than relying solely on narrative impressions. Anxiety in children and adolescents can be developmentally appropriate in some contexts but may become clinically significant when persistent, excessive or impairing. A family-centred consultation can clarify triggers and coping patterns. However, a homeopathic remedy should not be presented as a substitute for psychological therapy, school support or pharmacological treatment when these are indicated. The 2024 evidence evaluation included anxiety among its categories, illustrating how broad the research field has become without eliminating uncertainty about specific benefit (Maurin et al., 2024).

19.Organ-system symptoms that require a broader differential diagnosis:

One advantage of an anatomical framework is that it prevents symptoms from being treated as isolated complaints. Cough, for example, can arise from infection, asthma, upper-airway disease, aspiration, environmental exposure or cardiac disease. Vomiting can reflect infection, feeding intolerance, gastro-oesophageal disorders, obstruction, metabolic disease or intracranial pathology. Fatigue can be related to sleep, nutrition, anaemia, endocrine disorders, chronic inflammation or psychosocial stress. A homeopathic symptom picture may contain rich descriptive detail, but the differential diagnosis still depends on physiology and clinical examination. The same principle applies to recurrent fever. A child who has frequent mild viral infections in daycare may have a very different explanation from a child with prolonged fever, poor growth, unusual infections or inflammatory features. Before attributing recurrent symptoms to an individual “susceptibility”, clinicians should review the number and severity of episodes, microbiological diagnoses when available, exposure patterns, vaccination status, growth, family history and relevant laboratory evaluation. Such a framework reduces the risk of turning normal childhood exposure into an inappropriate disease label. Symptoms involving the nervous system deserve particular caution because behavioural and neurological signs can be subtle. A child who appears inattentive may be tired, anxious, have hearing

or vision problems, have a learning disorder, have ADHD, or simply be responding to an unsuitable learning environment. A child who develops a new movement abnormality, seizure, progressive weakness, regression of skills or persistent morning headache requires neurological assessment. The detailed history encouraged by individualised consultation can be helpful, but it should strengthen referral decisions rather than delay them. Cardiorespiratory findings also require close attention to severity. A "quiet" infant who is feeding poorly may have early respiratory compromise even before dramatic signs appear. Tachypnoea can accompany fever, metabolic acidosis or pulmonary disease. Cyanosis can have cardiac, respiratory or circulatory causes. When vital signs or appearance are abnormal, the child's physiological reserve should guide the urgency of evaluation. No complementary framework should override objective evidence of deterioration.

20.Special considerations in neonates, children with chronic disease and adolescents:

Neonates represent a special population because physiological adaptation is still taking place. Temperature control, glucose regulation, respiratory transition, feeding coordination and fluid balance can change rapidly. Symptoms may be nonspecific; poor feeding, lethargy, temperature instability or a change in breathing pattern can be the earliest signs of serious illness. Because evidence for complementary therapies is particularly limited in neonates, the threshold for specialist paediatric assessment should be low.

Children with chronic conditions also require a more complex medication and organ-function review. Examples include congenital heart disease, chronic lung disease, epilepsy, diabetes, kidney disease and neurodevelopmental disorders. These children may take several medicines, increasing the possibility of interactions and care fragmentation. Developmental pharmacology demonstrates that organ maturation changes medicine disposition, while disease itself may alter clearance and distribution. For this reason, complementary products should be disclosed to the prescribing team and included in medication reconciliation (van den Anker et al., 2018; Chapron et al., 2022). Adolescents require consideration of autonomy, privacy and mental health. Confidential conversation may reveal self-harm risk, substance use, disordered eating, sexual-health concerns or family conflict that caregivers do not initially report. A homeopathic consultation focused only on temperament or constitutional themes can miss these risks if it does not include standard adolescent health screening. The continuing maturation of executive and social brain networks also means that adolescents may experience symptoms differently from younger children (Dow-Edwards et al., 2019).

21.Clinical documentation and outcome tracking:

Good documentation makes complementary care safer and more scientifically interpretable. A paediatric record should identify the presenting diagnosis or working diagnosis, date of onset, age, weight and relevant developmental information, examination findings, current conventional treatment, complementary products already being used, red flags screened, caregiver expectations, and the agreed follow-up plan. The record should also state whether the complementary intervention is adjunctive and what conventional treatment remains necessary.

Outcome tracking should use both symptom and function. For a child with eczema, this may include itch severity, sleep disruption, affected body surface and validated quality-of-life measures. For recurrent infections, confirmed episode count, antibiotic exposure, missed school days and complications are more informative than a general statement that the immune system seems "better". In respiratory complaints, symptom-free days, exacerbations and acute-care utilisation are preferable to a simple subjective impression. The choice of outcome should be made before treatment whenever possible. Repeated measurement also helps clinicians recognise regression to the mean and natural fluctuation. Chronic paediatric conditions often have variable periods of worsening and improvement. Starting treatment during a symptom peak can be followed by improvement even when the intervention has no specific effect. Predefined follow-up intervals and objective measures reduce this interpretive problem. For research, the same principle supports more reliable comparisons and reduces selective reporting.

ISSN: 2584-1874; SJBLS
Scienxt Journal of Biotechnology and Life Science

22.Developmentally appropriate communication during individualised care:

Communication in paediatrics occurs with the child and caregiver, not only with the person who makes the appointment. Infants and toddlers communicate through behaviour, feeding and physiologic signs; school-aged children can usually provide more specific symptom information; adolescents may prefer direct discussion. A clinician should therefore adapt language, observe non-verbal responses and avoid suggestive questioning that encourages the child to endorse a particular narrative.

Caregivers should be asked what they believe the symptom means and what outcome they hope to achieve. This creates an opportunity to correct misconceptions without confrontation. For example, recurrent viral infections can be explained in terms of normal exposure and maturation rather than automatically describing the child as immunologically weak. Families can also be told why evidence-based treatments are recommended and how complementary care can or cannot be combined with them.

Such conversations preserve autonomy while protecting the child. The same communication principle applies when evidence is uncertain. A fair explanation can acknowledge that some studies report improvement while higher-quality evidence has not consistently demonstrated a specific effect. This is more informative than an absolute promise or a blanket dismissal. Parents can then decide what aligns with their values while remaining aware of the medical risks of delaying proven care.

Table.1: Summary table: developmental features that influence paediatric assessment

Domain	Developmental change	Practical relevance to complementary assessment
Body size and composition	Rapid changes in weight, body water and body fat across childhood	Record current weight and growth trajectory; do not assume adult physiology
Brain and behaviour	Ongoing synaptic refinement, myelination and executive development	Interpret behaviour by developmental stage; assess sleep, school and neurodevelopment
Airways and lungs	Smaller airways and continuing postnatal lung maturation	Screen carefully for respiratory distress; objective assessment takes priority
Kidneys and fluid balance	Maturation of filtration and tubular handling	Pay close attention to hydration and urine output during acute illness
Liver and drug disposition	Changing enzyme, transporter and binding capacity	Review all medicines and complementary products; age affects exposure
Immune system	Progressive maturation shaped by environment, vaccines and microbiome	Differentiate common exposure from unusual/recurrent severe infection
Endocrine/puberty	Rapid hormonal and body-composition transitions in adolescence	Interpret fatigue, weight change and menstrual symptoms in developmental context
Skin and thermoregulation	Developing barrier function and large surface-area-to-mass ratio in young children	Assess rash severity, infection, hydration and temperature regulation
Communication	Increasing capacity for history, consent and self-report with age	Adapt questioning; involve the child and respect adolescent confidentiality

23.Integrating Homeopathy with Evidence-Based Paediatric Care:

Integrative care is safest when it is additive rather than substitutive. This means that a child should receive appropriate diagnosis, preventive care, vaccination, nutrition advice, emergency recognition and proven treatment before a complementary approach is considered. The complementary intervention is then evaluated for whether it adds meaningful benefit without causing direct harm, interactions or delay. NCCIH guidance specifically advises parents not to replace or delay conventional care with complementary approaches that have not been shown to be safe and effective (NCCIH, 2025a).

Communication is central. Parents may have strong reasons for choosing homeopathy, including previous experiences, cultural beliefs, concerns about adverse effects or a preference for personalised care. Dismissing those concerns can weaken therapeutic relationships. A better approach is to ask what they hope the treatment will achieve, explain what is known about the evidence, identify safety issues, and make a shared plan for monitoring. The clinician should also document all complementary products so that possible interactions and duplicate therapies can be recognised.

The concept of "watchful waiting" should be used carefully. For a self-limiting viral illness in an otherwise well child, observation with supportive care may be appropriate. That is not evidence that a homeopathic remedy caused improvement. Conversely, when a serious disease is present, observation without effective therapy can create avoidable risk. Distinguishing these situations requires clinical diagnosis rather than the perceived gentleness of an intervention.

In multidisciplinary practice, pharmacists and nurses can contribute by reconciling all products, checking formulation and ingredient information, monitoring symptoms and adverse events, reinforcing vaccination and nutrition, and identifying deterioration. Paediatricians can provide diagnostic oversight and define escalation criteria. Complementary practitioners can contribute detailed symptom histories and support communication, provided that scope of practice and referral responsibilities are clear.




Figure.2: Evidence-informed integration of paediatric homeopathy with conventional care

24. Safety, Pharmacovigilance and Ethical Considerations:

Paediatric safety is broader than adverse effects from the product itself. It includes diagnostic error, treatment delay, interactions, contamination or incorrect labelling, excessive caregiver burden, financial harm and failure to escalate when symptoms worsen. Reviews of homeopathy-related

adverse events have documented both direct reactions and harm associated with replacing conventional care, while also noting limitations in the reporting literature (Posadzki et al., 2012). A more recent systematic review of observational studies found that adverse effects were reported in homeopathic research but emphasised the need for standardised reporting systems (Stub et al., 2022).
Product quality matters because the label "homeopathic" does not guarantee pharmacological inactivity. Some products may contain appreciable active ingredients or have manufacturing and dilution characteristics that differ from expectations (NCCIH, 2025b). For children, accidental ingestion is another practical concern. Products should be stored safely, and caregivers should keep accurate records of all medicines, supplements and complementary preparations.
The ethical principle of non-maleficence is particularly important in serious paediatric illness. A child with bacterial meningitis, sepsis, severe asthma, diabetic ketoacidosis, dehydration, anaphylaxis, significant trauma or suspected malignancy needs timely evidence-based care. Homeopathy may be discussed only in a way that preserves access to necessary treatment. The clinician should communicate clear warning signs and a specific follow-up plan.
Informed decision-making should also recognise the child's developing autonomy. Older children and adolescents should be included in explanations, and adolescents may need confidential discussion of sensitive topics. Caregivers should receive balanced information rather than promises of "natural", "side-effect-free" or "immune-boosting" treatment. Statements about efficacy should correspond to the strength of evidence rather than the number of families reporting benefit. WHO guidance on medicine safety emphasises the particular importance of monitoring medicines in children because evidence generated in adults cannot simply be transferred to younger populations (World Health Organization, 2007). The same principle supports stronger pharmacovigilance for complementary products. Every paediatric service that permits or records complementary use should have a process for identifying adverse events, documenting suspected products and communicating concerns to the relevant clinician or regulatory authority.

25. Evidence Appraisal: What the Current Literature Can and Cannot Tell Us:

The evidence base for homeopathy illustrates a recurring methodological problem in complementary medicine: studies may test different interventions while using the same label. An individualised homeopathic consultation is not identical to a fixed over-the-counter preparation. Some trials emphasise symptom outcomes; others focus on quality of life, recurrence, healthcare use or laboratory markers. This heterogeneity makes it difficult to combine studies and can produce apparently conflicting results.
Early systematic review work in paediatric homeopathy found mixed findings for a small number of conditions and insufficient evidence for reliable recommendations (Altunç et al., 2007). Broader comparative analyses have raised concerns that smaller and methodologically weaker studies can generate more favourable estimates than larger, more rigorous trials (Shang et al., 2005). More recent reviews continue to report uncertainty across specific paediatric conditions, with higher-quality respiratory trials failing to demonstrate a consistent specific benefit (King et al., 2021; Hawke et al., 2022). The 2024 Australian evidence evaluation is particularly useful because it updated the evidence base across many clinical categories using a structured methodology. It reviewed atopic conditions, recurrent childhood infections, neurodevelopmental disorders, asthma, diarrhoea, infantile colic, anxiety and other conditions. The report was prepared by HTANALYSTS and approved by the NHMRC Natural Therapies Working Committee in August 2024 (Maurin et al., 2024). Its conclusions provide an important contemporary benchmark for interpreting the paediatric literature. Evidence should also be separated into effectiveness and experience. Parents may genuinely observe improvement after beginning homeopathy. That observation deserves respectful listening, but the cause of improvement can have several explanations: natural history, regression to the mean, improved routine, increased caregiver attention, concurrent treatment, placebo or contextual effects, or a specific effect of the homeopathic product. Only appropriate comparative research can distinguish these possibilities reliably. This is why good clinical practice can include empathy and individualisation while still maintaining evidentiary discipline.

Table.2: Evidence Appraisal

Evidence feature	Why it matters in paediatric homeopathy	Preferred research practice
Randomisation and allocation concealment	Reduces systematic differences between treatment groups	Use computer-generated randomisation, concealment and prespecified analysis
Blinding	Limits expectation effects in subjective outcomes	Blind caregivers, participants and assessors where feasible; describe any unavoidable unblinding
Transparent individualisation	Homeopathic remedy selection may vary between children	Publish consultation protocol, repertory method, remedy decision rules and practitioner training
Age stratification	Development changes symptoms, physiology and treatment response	Report outcomes by age/developmental band and relevant maturation variables
Validated outcomes	Subjective improvement can overestimate specific treatment effects	Use validated symptom, function, quality-of-life and developmental measures
Treatment displacement	Harm may occur through delayed standard care	Record all conventional treatments and any care withheld or delayed
Adverse-event reporting	Children may experience different risks from adults	Use standard definitions, severity grades, product details and follow-up duration
Replication	Single positive trials may be unstable	Use multicentre replication and independent analysis

26.Research Priorities for a Developmentally Informed Evidence Base:

Future research should move beyond small single-centre trials that provide limited information about how the intervention was delivered. A robust trial should define the target condition using accepted paediatric diagnostic criteria, prespecify primary and secondary outcomes, report age and developmental stage, and describe conventional care received by every participant. When individualisation is central to the intervention, the consultation process should be protocolised without eliminating clinically meaningful flexibility. Age is a particularly important variable. A remedy or consultation may appear different in infants, preschool children and adolescents simply because the symptom profile and natural history differ. Investigators should therefore avoid treating a wide age range as a single homogeneous population. Growth, pubertal status, neurodevelopmental profile and baseline severity may also modify outcomes. Developmental pharmacology provides a model for this type of thinking because age-dependent physiology is treated as a measurable source of variation rather than noise (Chapron et al., 2022). Outcome selection should reflect what matters to children and families. Symptom intensity alone may not capture clinically meaningful change. Examples include fewer school absences, better sleep, reduced rescue-medication use, improved exercise tolerance, improved eczema-related quality of life, reduction in crying time, fewer medically confirmed infections, or stable growth. Objective measurements should be used when available, and caregiver-reported outcomes should be interpreted with awareness of expectation effects.

Pragmatic trials can complement placebo-controlled trials by testing homeopathy as actually delivered in practice, including consultation time and follow-up. However, pragmatic designs still require careful comparator definition and transparent reporting. A trial comparing a full homeopathic service with usual care answers a different question from a double-blind remedy-versus-placebo trial.

Both can be informative when their research questions are explicit. Longitudinal observational studies may help describe patterns of use and safety, but they should not be treated as proof of efficacy. Registries should capture the reason for choosing homeopathy, the clinical diagnosis, products used, conventional treatment, adverse events, and whether any effective treatment was delayed. Such data would improve understanding of real-world practice and help identify safety signals that randomised trials may be too small to detect. Mechanistic research also requires caution. A biological observation, such as a change in a biomarker after a consultation, does not automatically show that the homeopathic preparation caused the change. Studies should include appropriate controls, dose or dilution characterisation, assay validation and independent replication. Claims about memory of water, specific energetic signatures or direct organ-system modulation should be considered hypotheses until supported by reproducible, clinically relevant evidence.




Figure.3: Safety and research-governance framework for paediatric homeopathy

27.Practical Clinical Principles for Paediatric Homeopathic Settings:

A developmentally informed paediatric homeopathic setting should function first as a safe clinical environment. Basic observations such as weight, growth trajectory, temperature and respiratory status can identify children who need escalation. The consultation should document past diagnoses and treatments, including medicines prescribed by other clinicians. Families should be encouraged to maintain a single, current medication list rather than assuming that complementary products can be kept separate from conventional care.

The second principle is diagnostic humility. Common symptoms such as cough, diarrhoea, headache, abdominal pain, rash or tiredness have broad differentials. A characteristic symptom pattern may be useful for communication, but it does not narrow the differential sufficiently by itself. When symptoms are persistent, severe, recurrent or atypical, the clinician should seek appropriate diagnostic evaluation. The third principle is measurable follow-up. A treatment plan should state what improvement would look like, how it will be measured, and when the child will be reviewed. For example, improvement could be defined as reduced symptom frequency, better sleep, stable growth or improved activity. The plan should also identify what counts as non-response and which symptoms require urgent reassessment.

The fourth principle is transparent communication about evidence. Rather than telling families that homeopathy "works" or "does not work" in an undifferentiated way, clinicians can explain that evidence varies by condition and that higher-quality studies have not established a consistent specific effect for common paediatric indications. This approach allows families to make informed choices while maintaining trust and avoiding overstatement. The fifth principle is protection from treatment displacement. Complementary care is safest when it is used alongside, rather than instead of, proven interventions. Vaccinations, oral rehydration, antibiotics when clearly indicated, inhaled therapies for

asthma, emergency treatment for anaphylaxis and other established interventions should not be withheld because of reliance on homeopathy. The ethical objective is not to win an argument about medical philosophy; it is to preserve the child's access to timely and effective care.

28. Discussion:

The anatomical and physiological perspective changes how paediatric homeopathy should be interpreted. The strongest scientific foundation in the topic is the developmental biology of the child: organs mature at different rates, symptoms change with age, and treatment risk depends on physiology and severity. Homeopathy adds a distinctive individualised consultation model, but its proposed therapeutic effects must be evaluated independently from the valuable components of careful history taking, observation and caregiver engagement.

This distinction also helps explain why parents can report meaningful benefit even when controlled trials do not show a specific effect. A long consultation may reveal sleep problems, feeding difficulties, environmental triggers or family stressors that were previously overlooked. Once these factors are addressed, the child may improve. Such improvement is clinically real, but it does not identify which component of care caused it. Research needs to disentangle consultation effects, co-interventions and remedy effects without dismissing the lived experience of families. Organ-system development is particularly important in respiratory, gastrointestinal, neurological and immune complaints. The smaller paediatric airway, changing fluid balance, evolving brain networks and developing immune system all create age-specific clinical contexts. A treatment that seems "gentle" cannot be evaluated solely by the absence of an obvious adverse effect. The relevant safety question is whether the child received the correct diagnosis and appropriate treatment at the right time.

The current evidence also illustrates the importance of independent replication. A field can accumulate many positive small trials without producing a stable clinical conclusion. The contrast between some earlier positive findings and more recent higher-quality evidence in paediatric respiratory infections is an example of why trial quality and replication matter (King et al., 2021; Hawke et al., 2022). The 2024 Australian evidence evaluation provides an updated multi-condition framework but does not eliminate the underlying heterogeneity of the primary literature (Maurin et al., 2024).

A balanced interpretation therefore recognises three simultaneous facts. First, individualised care, careful listening and shared decision-making are legitimate components of child-centred practice. Second, homeopathic efficacy for paediatric conditions remains unproven or uncertain across much of the literature, and higher-quality studies have often failed to demonstrate specific benefit. Third, complementary use is not automatically harmless, particularly when products contain active ingredients or when effective treatment is delayed. These points can coexist without requiring an all-or-none position.

29. Conclusion:

Paediatric homeopathy can only be responsibly understood when the biology of the developing child remains central. Growth, organ maturation, neural development, respiratory mechanics, immune education, gastrointestinal ecology, renal clearance, endocrine change and changing behaviour all influence symptom expression and clinical risk. The same symptom has different meaning across developmental stages, and any individualised consultation must account for these differences.

The evidence reviewed does not establish that homeopathic preparations reliably modify paediatric organ-system physiology or produce consistent specific benefits across common childhood conditions. Reviews of respiratory infections, eczema, childhood and adolescent disorders, and more recent government evidence evaluations indicate continuing uncertainty and frequent methodological limitations. Some individual studies report positive outcomes, but these findings should be interpreted within the broader body of evidence rather than in isolation.

The most defensible clinical role for homeopathy in paediatric settings, when families choose to use it, is therefore as a carefully monitored complementary approach that does not replace diagnosis, prevention, emergency care or treatments with established benefit. Future studies should be

developmentally informed, transparent about individualisation, rigorous in outcome measurement, and explicit about adverse events and treatment displacement. Such an approach respects both the interest in personalised care and the obligation to protect children through scientifically accountable practice.

30. References:

(1) World Health Organization. WHO child growth standards: length/height-for-age, weight-for-age, weight-for-length/height, weight-for-height and body mass index-for-age: methods and development. Geneva: World Health Organization; 2006.
(2) World Health Organization. Child growth standards [Internet]. Geneva: World Health Organization; 2025 Feb 12 [cited 2026 Oct 1]. Available from: https://www.who.int/news-room/questions-and-answers/item/child-growth-standards
(3) Kearns GL, Abdel-Rahman SM, Alander SW, Blowey DL, Leeder JS, Kauffman RE. Developmental pharmacology-drug disposition, action, and therapy in infants and children. N Engl J Med. 2003;349(12):1157-67. doi:10.1056/NEJMra035092.
(4) van den Anker J, Reed MD, Allegaert K, Kearns GL. Developmental changes in pharmacokinetics and pharmacodynamics. J Clin Pharmacol. 2018;58 Suppl 10:S10-S25. doi:10.1002/jcph.1284.
(5) Russel FG, et al. Physiologically-based pharmacokinetic models for children: starting to reach maturation? Pharmacol Ther. 2020;211:107541. doi:10.1016/j.pharmthera.2020.107541.
(6) Chapron BD, Chapron A, Leeder JS. Recent advances in the ontogeny of drug disposition. Br J Clin Pharmacol. 2022;88(10):4267-84. doi:10.1111/bcp.14821.
(7) Mullassery D, Smith NP. Lung development. Semin Pediatr Surg. 2015;24(4):152-5. doi:10.1053/j.sempedsurg.2015.01.011.
(8) Stocks J. The functional growth and development of the lung during the first year of life. Early Hum Dev. 1977;1(3):285-309. doi:10.1016/0378-3782(77)90041-X.
(9) Stocks J, Hislop A, Sonnappa S. Early lung development: lifelong effect on respiratory health and disease. Lancet Respir Med. 2013;1(9):728-42. doi:10.1016/S2213-2600(13)70118-8.
(10) Pelizzo GP, Calcaterra VC, Baldassarre P, Marinaro M, Taranto S, Ceresola M, et al. The impact of hormones on lung development and function: an overlooked aspect to consider from early childhood. Front Endocrinol (Lausanne). 2024;15:1425149. doi:10.3389/fendo.2024.1425149.
(11) Taki Y, Kawashima R. Brain development in childhood. Open Neuroimag J. 2012;6:103-10. doi:10.2174/1874440001206010103.
(12) Dow-Edwards D, MacMaster FP, Peterson BS, Niesink R, Andersen S, Braams BR. Experience during adolescence shapes brain development: from synapses and networks to normal and pathological behavior. Neurotoxicol Teratol. 2019;76:106834. doi:10.1016/j.ntt.2019.106834.
(13) Djuardi Y, Wammes LJ, Supali T, Sartono E, Yazdanbakhsh M. Immunological footprint: the development of a child's immune system in environments rich in microorganisms and parasites. Parasitology. 2011;138(12):1508-18. doi:10.1017/S0031182011000588.
(14) Li M, Wang M, Donovan SM. Early development of the gut microbiome and immune-mediated childhood disorders. Semin Reprod Med. 2014;32(1):74-86. doi:10.1055/s-0033-1361825.
(15) Ahearn-Ford S, Berrington JE, Stewart CJ. Development of the gut microbiome in early life. Exp Physiol. 2022;107(5):415-21. doi:10.1113/EP089919.
(16) World Health Organization. Promoting safety of medicines for children. Geneva: World Health Organization; 2007.
(17) National Center for Complementary and Integrative Health. Children and the use of complementary health approaches [Internet]. Bethesda (MD): NCCIH; 2025 [cited 2026 Oct 1]. Available from: https://www.nccih.nih.gov/health/children-and-the-use-of-complementary-health-approaches

(18) National Center for Complementary and Integrative Health. Homeopathy: what you need to know [Internet]. Bethesda (MD): NCCIH; 2025 [cited 2026 Oct 1]. Available from: https://www.nccih.nih.gov/health/homeopathy
(19) Hawke K, King D, van Driel ML, McGuire TM. Homeopathic medicinal products for preventing and treating acute respiratory tract infections in children. Cochrane Database Syst Rev. 2022;12(12):CD005974. doi:10.1002/14651858.CD005974.pub6.
(20) King D, Hawke K, McGuire TM, van Driel M. Homeopathic preparations for preventing and treating acute upper respiratory tract infections in children: a systematic review and meta-analysis. Acad Pediatr. 2021;21(2):211-20. doi:10.1016/j.acap.2020.07.016.
(21) Altunç U, Pittler MH, Ernst E. Homeopathy for childhood and adolescence ailments: systematic review of randomized clinical trials. Mayo Clin Proc. 2007;82(1):69-75. doi:10.4065/82.1.69.
(22) Banerjee K, Mathie RT, Costelloe C, Howick J. Homeopathy for allergic rhinitis: a systematic review. J Altern Complement Med. 2017;23(6):426-44. doi:10.1089/acm.2016.0310.
(23) Ernst E. Homeopathy for eczema: a systematic review of controlled clinical trials. Br J Dermatol. 2012;166(6):1170-2. doi:10.1111/j.1365-2133.2012.10994.x.
(24) Shang A, Huwiler-Müntener K, Nartey L, Jüni P, Dörig S, Sterne JAC, et al. Are the clinical effects of homoeopathy placebo effects? Comparative study of placebo-controlled trials of homoeopathy and allopathy. Lancet. 2005;366(9487):726-32. doi:10.1016/S0140-6736(05)67177-2.
(25) Maurin R, Rogers E, Sullivan E, Nolan K, Denham I, Jorgensen MA. Homeopathy for preventing and treating health conditions: evidence evaluation report. Canberra: National Health and Medical Research Council; 2024.