

Pain management techniques in medical-surgical nursing: Exploring pharmacological and non-pharmacological methods

Manmeet Kaur^{1*}, Lingaraju AR², Shrushti Bandbe³

^{1*}Associate Professor, University College of Nursing Faridkot, Punjab, India

²Professor, Department of Community Health Nursing, MES College of Nursing, Ghanekhunt-Lote, Ratnagiri, Maharashtra, Affiliated by MUHS, Nashik, India

³Assistant Professor, Department of Community Health Nursing, MES College of Nursing, Ghanekhunt-Lote, Ratnagiri, Maharashtra, Affiliated by MUHS, Nashik, India

<https://doi.org/10.5281/zenodo.19688995>

Abstract:

Pain management is a fundamental aspect of medical-surgical nursing that directly influences patient recovery, satisfaction, and overall quality of care. Effective pain control requires a comprehensive and evidence-based approach that integrates both pharmacological and non-pharmacological strategies tailored to individual patient needs. Pharmacological methods, including non-opioid analgesics, opioids, and adjuvant medications, remain essential for managing varying intensities and types of pain, particularly in acute and postoperative conditions. However, increasing concerns regarding adverse effects and opioid dependence have highlighted the importance of incorporating non-pharmacological interventions such as cognitive-behavioral techniques, physical therapies, and complementary approaches like relaxation, music therapy, and transcutaneous electrical nerve stimulation. Nurses play a pivotal role in pain management through continuous assessment, timely intervention, patient education, and evaluation of outcomes, ensuring a patient-centered and holistic approach to care. The integration of multimodal pain management strategies has been shown to enhance analgesic effectiveness, reduce medication-related complications, and improve patient comfort. Furthermore, emerging technologies such as telehealth and artificial intelligence are transforming pain assessment and management practices.

Keywords:

Pain Management, Medical-Surgical Nursing, Pharmacological Methods, Non-Pharmacological Interventions, Multimodal Approach, Nursing Care, Patient-Centered Care, Analgesics.

1. Introduction:

Pain is widely recognized as a complex and multidimensional experience that encompasses not only physiological mechanisms but also psychological and social influences, making it a highly individualized phenomenon that varies across patients and clinical contexts, and contemporary frameworks such as the biopsychosocial model emphasize that pain arises from a dynamic interaction between biological processes, emotional responses, and social environments, thereby challenging traditional biomedical perspectives that focus solely on tissue damage (Smith et al., 2021). This understanding highlights that pain is not merely a sensory event but also an emotional and cognitive experience shaped by prior experiences, beliefs, and expectations, which explains why individuals respond differently to similar painful stimuli (Johnson et al., 2019). In medical-surgical settings, pain is particularly prevalent due to surgical trauma, inflammatory responses, and underlying disease conditions, and it may manifest as acute pain that serves a protective function or chronic pain that persists beyond healing and contributes to long-term disability and reduced quality of life (Williams et al., 2020). Moreover, psychological factors such as anxiety, depression, fear, and catastrophizing can amplify pain perception and interfere with recovery, while social determinants including cultural beliefs, family support, and socioeconomic status further influence how pain is perceived, expressed, and managed (Brown et al., 2018). These multidimensional influences significantly affect patient outcomes, as poorly managed pain is associated with delayed wound healing, prolonged hospitalization, increased healthcare costs, and diminished patient satisfaction, thereby underscoring the need for comprehensive

and individualized pain management strategies (Taylor et al., 2022). Within this context, nurses play a pivotal role in pain management as they are continuously present at the bedside, enabling them to perform ongoing assessment, monitor treatment effectiveness, and implement timely interventions that address both physical and psychosocial aspects of pain (Anderson et al., 2021). Their responsibilities extend beyond medication administration to include patient education, emotional support, and advocacy, ensuring that pain is recognized, reported, and managed appropriately according to evidence-based guidelines (Clark et al., 2019). Furthermore, the integration of nursing interventions within a holistic framework allows for the identification of factors that may exacerbate or alleviate pain, such as environmental stressors, patient beliefs, and coping mechanisms, thereby enhancing the effectiveness of care (Lewis et al., 2020). In addition, the variability in pain experiences among patients necessitates the use of individualized care plans that incorporate both pharmacological and non-pharmacological approaches, reflecting the understanding that no single intervention is universally effective for all patients (Walker et al., 2017). The importance of this comprehensive approach is reinforced by growing evidence suggesting that addressing psychological and social dimensions alongside physical symptoms leads to improved functional outcomes, greater patient satisfaction, and more efficient use of healthcare resources (Hall et al., 2023). Consequently, modern medical-surgical nursing practice emphasizes the need for a patient-centered and multidisciplinary approach to pain management, where nurses collaborate with other healthcare professionals to deliver coordinated and holistic care (Martin et al., 2020). This approach is particularly relevant in the context of increasing recognition of chronic pain as a major public health issue, requiring long-term management strategies that go beyond symptom control to address underlying contributing factors (Evans et al., 2018). Ultimately, the multidimensional nature of pain and the integrative role of nursing interventions, as illustrated in Figure 1, highlight the necessity of adopting a comprehensive and evidence-based approach to pain management that considers the full spectrum of factors influencing patient experience and recovery (Roberts et al., 2022).




Figure. 1: Conceptual framework of pain management in medical-surgical nursing illustrating biological, psychological, and social dimensions of pain

2. Physiology and Classification of Pain:

Pain is broadly classified into nociceptive, neuropathic, and mixed types, each representing distinct underlying mechanisms and clinical characteristics that require tailored management strategies in medical-surgical nursing practice, as nociceptive pain arises from the activation of peripheral nociceptors in response to tissue injury or inflammation and typically presents as sharp, aching, or throbbing sensations associated with identifiable damage (Kataria et al., 2024). Neuropathic pain, in contrast, results from lesions or diseases affecting the somatosensory nervous system and is often described as burning, tingling, or electric shock-like sensations, reflecting abnormal

neural processing rather than direct tissue injury (Leone et al., 2024). This distinction is clinically significant because neuropathic pain often involves central and peripheral sensitization mechanisms that alter neuronal excitability and reduce responsiveness to conventional analgesics such as opioids (Rugnath et al., 2024). Furthermore, nociceptive pain is considered a protective physiological response that alerts the body to potential harm, whereas neuropathic pain is typically maladaptive and persists even in the absence of ongoing tissue damage, contributing to chronic pain conditions and reduced quality of life (Pessôa et al., 2024). In many clinical cases, patients present with mixed pain, where both nociceptive and neuropathic components coexist, making assessment and management more complex and requiring a multimodal therapeutic approach (Wiesenfeld-Hallin et al., 2022). The physiological pathways involved in pain perception begin with the transduction of noxious stimuli into electrical signals at peripheral nociceptors, followed by transmission through afferent nerve fibers to the spinal cord and brain, where modulation and perception occur (Pacifico et al., 2024). These processes involve intricate neurochemical interactions, including neurotransmitters and ion channels that regulate pain signaling and contribute to sensitization phenomena in both acute and chronic pain states (Anderson et al., 2023). Central sensitization, in particular, plays a critical role in amplifying pain signals within the central nervous system, leading to heightened sensitivity and persistent pain even after the initial injury has resolved (Rugnath et al., 2024). Peripheral sensitization, on the other hand, occurs at the site of injury where inflammatory mediators lower the threshold of nociceptors, resulting in increased responsiveness to stimuli (Kataria et al., 2024). These mechanisms highlight the importance of understanding the underlying pathophysiology of pain in order to implement effective and targeted interventions that address both the source and perception of pain (Leone et al., 2024). For nurses, this knowledge is essential in guiding clinical decision-making, particularly in selecting appropriate pharmacological treatments such as NSAIDs for nociceptive pain and anticonvulsants or antidepressants for neuropathic pain (Pessôa et al., 2024). Additionally, recognizing the presence of mixed pain conditions allows nurses to integrate non-pharmacological strategies alongside medications, thereby enhancing overall pain relief and minimizing adverse effects (Kataria et al., 2024). The complexity of pain mechanisms also underscores the need for comprehensive assessment tools that can differentiate between pain types and guide individualized care plans (Leone et al., 2024). Moreover, advances in neurophysiological testing and imaging techniques have improved the understanding of pain pathways and contributed to more accurate diagnosis and targeted treatment approaches in clinical practice (Garcia et al., 2021). The integration of this knowledge into nursing practice enhances the ability to monitor patient responses, evaluate treatment effectiveness, and adjust interventions accordingly to achieve optimal outcomes (Rugnath et al., 2024). The physiological pathways involved in pain perception and classification are depicted in Figure 2, which demonstrates peripheral and central transmission mechanisms along with different categories of pain, thereby providing a visual representation of the complex processes underlying pain experience and reinforcing the importance of a mechanism-based approach to pain management in medical-surgical nursing (Kataria et al., 2024).




Figure. 2: Pain transmission pathways and classification showing nociceptive, neuropathic, and central mechanisms

3. Pain Assessment in Medical-Surgical Nursing:

Accurate pain assessment is essential for effective pain management, as pain is inherently subjective and varies significantly among individuals, making it necessary for healthcare professionals, particularly nurses, to rely on standardized and validated tools that can translate subjective experiences into measurable data to guide clinical decision-making and improve patient outcomes (Stretanski et al., 2025). The subjective nature of pain means that it cannot be directly observed or measured through objective parameters alone, and therefore patient self-report remains the gold standard in clinical practice, requiring tools that are simple, reliable, and adaptable across diverse patient populations (Hawker et al., 2017). Nurses utilize a range of standardized pain assessment tools, including the Visual Analog Scale, Numeric Rating Scale, and Faces Pain Scale, which are designed to quantify pain intensity while accommodating differences in age, cognitive ability, and communication capacity among patients (Breivik et al., 2018). The Numeric Rating Scale is widely used due to its simplicity, allowing patients to rate their pain on a scale from zero to ten, making it particularly effective in acute care settings where rapid assessment is required (Jensen et al., 2019). The Visual Analog Scale provides a continuous measure of pain intensity using a linear format, enabling more precise evaluation and monitoring of changes over time, which is especially useful in research and postoperative care settings (Hjermstad et al., 2018). The Faces Pain Scale is particularly beneficial for pediatric patients and individuals with communication difficulties, as it uses visual facial expressions to represent different levels of pain intensity, thereby enhancing patient understanding and participation in assessment (Tomlinson et al., 2019). The use of these tools is crucial in ensuring consistency in pain assessment, facilitating communication between patients and healthcare providers, and enabling the evaluation of treatment effectiveness over time (Karcioglu et al., 2018). The commonly used pain assessment tools and their clinical applications are illustrated in Figure 3, which highlights variations in tool selection based on patient characteristics such as age, cognitive status, and level of consciousness, thereby emphasizing the importance of individualized assessment approaches in nursing practice. In addition, the selection of appropriate pain assessment tools depends on multiple factors including patient literacy, cultural background, and clinical condition, as certain tools may not be suitable for all patients, particularly those with cognitive impairments or language barriers (Herr et al., 2019).




Figure. 3: Common pain assessment tools used in clinical practice across different patient populations

The effectiveness of pain assessment is further enhanced by combining multiple tools and approaches, including observational methods for non-verbal patients, which allow nurses to assess behavioral and physiological indicators of pain such as facial expressions, body movements, and vital signs (Chanques et al., 2017). The integration of these assessment methods ensures a more comprehensive understanding of the patient's pain experience and supports the development of individualized care plans that address both the intensity and underlying causes of pain (Pasero et al., 2018). Furthermore, ongoing pain assessment is essential for monitoring the effectiveness of interventions and making necessary adjustments to treatment plans, thereby improving patient outcomes and preventing complications associated with unmanaged pain (Gordon et al., 2020). The selection and characteristics of these tools are further summarized in Table 1, where Table 1 presents common pain assessment

tools, their clinical applications, advantages, and limitations within different patient populations, providing a structured overview that supports clinical decision-making and enhances the quality of pain management in medical-surgical nursing practice.

Table 1: Common pain assessment tools, their clinical applications, advantages, and limitations

Tool	Patient Type	Advantages	Limitations
VAS	Adults	Sensitive	Requires understanding
NRS	Adults	Easy to use	Subjective bias
Faces Scale	Children	Visual aid	Less precise

4. Pharmacological pain management:

Pharmacological interventions remain the cornerstone of pain management in medical-surgical nursing, particularly for patients experiencing moderate to severe pain, as these approaches provide rapid and effective relief through well-established mechanisms targeting nociceptive and neuropathic pathways (Chou et al., 2016). The selection of appropriate pharmacological agents depends on multiple factors including pain intensity, underlying etiology, patient comorbidities, and risk of adverse effects, requiring nurses to apply clinical judgment and evidence-based guidelines to ensure safe and effective treatment (Buvanendran et al., 2019). The World Health Organization analgesic ladder, widely recognized as a foundational framework for pain management, outlines a stepwise approach that begins with non-opioid medications for mild pain and progresses to stronger opioids for more severe pain conditions, as illustrated in Figure 4, thereby promoting rational and structured analgesic use (WHO et al., 2019). Non-opioid analgesics such as nonsteroidal anti-inflammatory drugs and acetaminophen are commonly used as first-line agents due to their effectiveness in reducing inflammation and peripheral sensitization, making them suitable for mild to moderate pain conditions (Moore et al., 2018). These medications are generally well tolerated but may be associated with adverse effects such as gastrointestinal irritation, renal impairment, and cardiovascular risks, particularly with prolonged use, necessitating careful monitoring by nurses (Derry et al., 2017).




Figure 4: WHO analgesic ladder demonstrating stepwise pharmacological pain management

Opioid analgesics, including morphine, fentanyl, and oxycodone, are indicated for moderate to severe pain and act primarily on central nervous system receptors to alter pain perception and response, providing potent analgesic effects (Volkow et al., 2019). Despite their effectiveness, opioids carry significant risks such as respiratory depression, sedation, tolerance, dependence, and potential for misuse, which require vigilant assessment, dose titration, and monitoring by nursing professionals (Dowell et al., 2022). Adjuvant medications, including antidepressants, anticonvulsants, corticosteroids, and muscle relaxants, are increasingly used as part of multimodal analgesia, particularly in neuropathic pain conditions where traditional analgesics may be insufficient (Finnerup et al., 2021). These agents target specific pain mechanisms such as nerve sensitization and central processing,

thereby enhancing analgesic outcomes and reducing reliance on opioids (Wiffen et al., 2020). The concept of multimodal pharmacological therapy emphasizes the use of multiple agents with different mechanisms of action to achieve synergistic effects, improve pain control, and minimize side effects associated with higher doses of single medications (Kehlet et al., 2018). Nurses play a critical role in administering these medications safely, monitoring therapeutic responses, identifying adverse effects, and educating patients about proper use and potential risks, thereby ensuring optimal outcomes (Gordon et al., 2020). The classification and clinical use of these pharmacological agents are summarized in Table 2, where Table 2 outlines different drug classes, examples, indications, and associated side effects relevant to nursing practice, providing a structured framework that supports clinical decision-making and enhances patient safety. Furthermore, individualized pharmacological management is essential, as patient-specific factors such as age, renal and hepatic function, and previous medication history influence drug selection and dosing (Pergolizzi et al., 2018). Advances in pharmacogenomics and personalized medicine are also contributing to more targeted pain management strategies by identifying genetic variations that affect drug metabolism and response (Smith et al., 2021). In addition, ongoing evaluation of pain relief and side effects is necessary to adjust treatment regimens and ensure sustained effectiveness, highlighting the dynamic nature of pharmacological pain management in clinical practice (Apfelbaum et al., 2018).

Table. 2: Pharmacological agents used in pain management including drug classes, examples, indications, and side effects

Drug Class	Examples	Indications	Side Effects
NSAIDs	Ibuprofen	Mild to moderate pain	Gastric irritation
Opioids	Morphine	Severe pain	Sedation
Adjuvants	Gabapentin	Neuropathic pain	Dizziness

5. Non-pharmacological pain management:

Non-pharmacological interventions are essential components of pain management that complement pharmacological therapies and enhance overall patient care by addressing the multidimensional nature of pain through physical, psychological, and emotional pathways, thereby improving patient comfort and reducing dependence on medications (Tick et al., 2018). These interventions are widely recognized for their cost-effectiveness, accessibility, and minimal side effects, making them particularly valuable in diverse clinical settings including medical-surgical units where patients may require continuous and supportive pain relief strategies (National Academies et al., 2019). The integration of non-pharmacological techniques into routine nursing care supports a holistic approach that not only alleviates pain but also promotes relaxation, improves coping mechanisms, and enhances overall patient well-being (Garland et al., 2020). The broad categories of non-pharmacological pain management techniques are illustrated in Figure 5, which classifies interventions into cognitive-behavioral, physical, emotional, and complementary approaches, thereby providing a structured framework for their application in clinical practice. Cognitive-behavioral strategies such as relaxation techniques, guided imagery, and distraction play a significant role in modifying pain perception by influencing the brain's interpretation of pain signals and reducing anxiety and stress associated with painful experiences (Eccleston et al., 2020). Relaxation techniques, including deep breathing and progressive muscle relaxation, help decrease sympathetic nervous system activity, leading to reduced muscle tension and improved pain tolerance (Zaccaro et al., 2018). Guided imagery allows patients to focus on positive mental images, thereby diverting attention away from pain and enhancing psychological resilience (Nguyen et al., 2019). Distraction techniques, such as engaging in conversation, listening to music, or watching television, are particularly effective in acute care settings where immediate pain relief is required (Birnie et al., 2017). Physical methods including massage, positioning, and heat or cold therapy contribute to pain relief by improving blood circulation, reducing inflammation, and alleviating muscle stiffness, making them essential components of nursing care for postoperative and musculoskeletal pain (Field et al., 2020). Massage therapy, for instance, has been shown to stimulate mechanoreceptors and promote the release of endorphins, thereby reducing pain intensity and enhancing relaxation (Li et al., 2021). Proper patient positioning is equally important in preventing pressure-related discomfort and improving functional mobility, particularly in bedridden patients (Kwon et al., 2019). Heat therapy helps in relaxing muscles and increasing blood flow, while cold therapy reduces inflammation and numbs affected areas, providing effective relief in different clinical scenarios (Malanga et al., 2015). Complementary therapies such as acupuncture, aromatherapy, and music

therapy further contribute to pain management by addressing both physiological and psychological aspects of pain, thereby promoting emotional well-being and patient satisfaction (Vickers et al., 2018).




Figure 5: Categories of non-pharmacological pain management techniques including cognitive, physical, and complementary approaches

Acupuncture, based on traditional Chinese medicine principles, involves the stimulation of specific points on the body to modulate pain pathways and restore balance, and has been widely supported by clinical evidence for its effectiveness in various pain conditions (Zhao et al., 2022). Music therapy has been shown to reduce anxiety, lower pain perception, and improve mood by influencing neurochemical pathways and providing a calming effect (Hole et al., 2015). Aromatherapy, using essential oils such as lavender, has demonstrated benefits in reducing stress and enhancing relaxation, thereby indirectly contributing to pain relief (Lakhan et al., 2016). The mechanisms and outcomes of these techniques are summarized in Table 3, where Table 3 highlights various non-pharmacological interventions, their mechanisms of action, and clinical outcomes in pain reduction and patient comfort, providing a comprehensive overview that supports evidence-based nursing practice. Furthermore, the effectiveness of non-pharmacological interventions is enhanced when they are tailored to individual patient preferences, cultural beliefs, and clinical conditions, ensuring a personalized approach to care that maximizes therapeutic benefits (Kwekkeboom et al., 2019). Nurses play a crucial role in implementing these interventions, educating patients about their use, and evaluating their effectiveness as part of a multimodal pain management strategy that integrates both pharmacological and non-pharmacological approaches (Gordon et al., 2020).

Table 3: Non-pharmacological pain management techniques with mechanisms of action and clinical outcomes

Technique	Mechanism	Outcome
Massage	Muscle relaxation	Pain reduction
CBT	Cognitive reframing	Improved coping
TENS	Nerve stimulation	Pain modulation
Music Therapy	Distraction	Anxiety reduction

6. Role of Nurses in Pain Management:

Nurses play a pivotal role in pain management within medical-surgical settings by conducting continuous and comprehensive assessments that ensure accurate identification of pain characteristics, intensity, and underlying causes, thereby forming the foundation for effective and individualized care planning (Gordon et al., 2020). Their role extends beyond initial assessment to include ongoing monitoring of patient responses to treatment, allowing timely modifications to interventions and ensuring optimal pain relief outcomes (Herr et al., 2019). Nurses are primarily responsible for administering pharmacological therapies safely, including non-opioid analgesics, opioids, and adjuvant medications, while carefully observing for therapeutic effects and potential adverse reactions to maintain patient safety (Pasero et al., 2018). In addition to medication administration, nurses implement a wide range of non-pharmacological interventions such as relaxation techniques, positioning, and distraction methods, which complement pharmacological approaches and contribute to holistic pain management (Kwekkeboom et al.,

2019). Patient education is another critical component of nursing responsibilities, as nurses provide information regarding pain management strategies, medication use, and self-care techniques, empowering patients to actively participate in their own care and improving adherence to treatment plans (Botti et al., 2018). Nurses also play a vital role in advocating for patients by ensuring that their pain is acknowledged, assessed regularly, and managed appropriately, particularly in situations where pain may be underestimated or undertreated due to communication barriers or clinical biases (Manias et al., 2020). Effective communication between nurses, patients, and multidisciplinary healthcare teams further enhances pain management by promoting coordinated care and ensuring that all aspects of the patient's condition are considered in decision-making processes (Booker et al., 2019). The integration of evidence-based practices into nursing care ensures that pain management strategies are aligned with current clinical guidelines and research findings, thereby improving patient outcomes and reducing the risk of complications associated with inadequate pain control (Tick et al., 2018). Nurses are also responsible for evaluating the effectiveness of pain management interventions through systematic reassessment and documentation, which supports continuity of care and facilitates quality improvement initiatives within healthcare settings (Chou et al., 2016). The comprehensive responsibilities of nurses in pain management are depicted in Figure 6, which illustrates their role across assessment, intervention, monitoring, and evaluation phases within clinical settings, highlighting the dynamic and multifaceted nature of nursing practice in pain management. Furthermore, nurses must consider individual patient factors such as age, cultural background, cognitive status, and comorbidities when developing and implementing pain management plans, ensuring that care is tailored to meet specific patient needs (Herr et al., 2019). The increasing emphasis on patient-centered care has further strengthened the role of nurses in involving patients and their families in decision-making processes, thereby enhancing satisfaction and improving overall healthcare experiences (Botti et al., 2018). Advances in healthcare technology, including electronic health records and digital pain assessment tools, have also supported nurses in documenting pain accurately and tracking patient progress over time, contributing to more efficient and effective care delivery (Manias et al., 2020). In addition, nurses play an essential role in identifying and addressing barriers to effective pain management, such as patient misconceptions, fear of addiction, and inadequate communication, thereby ensuring that patients receive appropriate and timely interventions (Booker et al., 2019). Their ability to combine clinical expertise with compassionate care enables them to address both the physical and emotional aspects of pain, ultimately improving patient comfort and quality of life (Tick et al., 2018).




Figure. 6: Nursing responsibilities in pain management including assessment, intervention, monitoring, and evaluation

7. Barriers to Effective Pain Management:

Despite significant advancements in pain management, multiple barriers continue to hinder its effective implementation in medical-surgical nursing practice, ultimately leading to suboptimal patient outcomes and compromised quality of care, as gaps in knowledge and training among healthcare professionals remain a persistent issue that affects accurate pain assessment and appropriate intervention selection (Alotaibi et al., 2021). Limited understanding of pain mechanisms and management strategies can result in underestimation of patient pain and

inadequate use of both pharmacological and non-pharmacological approaches, thereby reducing the effectiveness of care provided (Yava et al., 2018). Time constraints and heavy workloads further challenge nurses in busy clinical environments, where competing priorities may limit the opportunity for thorough pain assessment, documentation, and timely intervention (Kizza et al., 2019). In such settings, pain management may become secondary to other urgent clinical tasks, leading to delays in treatment and prolonged patient discomfort (Wells et al., 2020). Fear of opioid addiction and concerns about adverse effects represent another significant barrier, as both healthcare providers and patients may hesitate to use opioid medications even when clinically indicated, resulting in inadequate pain relief (Vadivelu et al., 2018). This fear is often compounded by the global opioid crisis, which has led to stricter prescribing practices and increased caution in opioid use, sometimes at the expense of effective pain control (Dowell et al., 2022). Cultural beliefs and attitudes toward pain also play a crucial role in influencing how patients perceive, express, and report pain, with some individuals underreporting pain due to cultural norms that discourage the expression of discomfort (Alhassan et al., 2020). Language barriers and differences in communication styles can further complicate pain assessment, making it difficult for nurses to accurately interpret patient needs and provide appropriate care (Samulowitz et al., 2018). Additionally, misconceptions about pain and its management among patients, such as the belief that pain is an inevitable part of illness or recovery, may prevent them from seeking timely relief and adhering to prescribed treatment plans (Gan et al., 2017). Healthcare provider-related factors, including inadequate adherence to clinical guidelines and variability in pain management practices, also contribute to inconsistencies in care delivery and patient outcomes (Meissner et al., 2018). System-related barriers such as limited resources, lack of standardized protocols, and insufficient staffing levels further impede the implementation of effective pain management strategies in healthcare settings (Gregory et al., 2019). The absence of institutional support and continuing education opportunities for healthcare professionals can hinder the adoption of evidence-based practices and limit the effectiveness of pain management interventions (Lui et al., 2021). Documentation issues and lack of proper pain assessment tools in some clinical settings may also result in incomplete or inaccurate recording of patient pain, affecting continuity of care and treatment planning (Booker et al., 2019). Psychological factors such as anxiety, depression, and fear experienced by patients can amplify pain perception and complicate management, requiring a more comprehensive and multidisciplinary approach to address both physical and emotional aspects of pain (Darnall et al., 2021). The various challenges associated with pain management are presented in Figure 7, which categorizes barriers into patient-related, healthcare provider-related, and system-related factors, thereby providing a structured understanding of the complexities involved in achieving effective pain control. Addressing these barriers requires a multifaceted approach that includes ongoing education and training for healthcare professionals, improved communication between patients and providers, and the development of standardized protocols that support consistent and evidence-based practice (Tick et al., 2018). Enhancing patient education and promoting awareness about pain management options can also empower individuals to actively participate in their care and improve treatment adherence (Gan et al., 2017). Furthermore, organizational support in the form of adequate staffing, access to resources, and implementation of quality improvement initiatives is essential for overcoming system-level challenges and ensuring effective pain management (Gregory et al., 2019). By recognizing and addressing these barriers, healthcare providers can improve pain management practices, enhance patient satisfaction, and achieve better clinical outcomes in medical-surgical nursing settings (Alotaibi et al., 2021).




Figure 7: Barriers to effective pain management categorized into patient-related, provider-related, and system-related factors

8. Multimodal pain management approach:

Multimodal pain management is a comprehensive approach that involves the integration of pharmacological and non-pharmacological strategies to achieve optimal pain control and improve overall patient outcomes in medical-surgical settings, as it targets multiple pain pathways simultaneously and provides synergistic effects that enhance analgesic efficacy (Kehlet et al., 2018). This approach is particularly effective in managing acute postoperative pain and chronic pain conditions, where reliance on a single modality may be insufficient to address the complex and multidimensional nature of pain (Buvanendran et al., 2019). By combining different classes of medications such as non-opioid analgesics, opioids, and adjuvant drugs with supportive non-pharmacological interventions, multimodal pain management reduces the need for high doses of individual drugs and minimizes the risk of adverse effects associated with monotherapy (Chou et al., 2016). The reduction in opioid consumption is one of the key benefits of this approach, as it helps decrease the likelihood of opioid-related complications such as respiratory depression, tolerance, and dependence, which are major concerns in modern clinical practice (Dowell et al., 2022). Non-pharmacological interventions such as relaxation techniques, physical therapy, cognitive-behavioral strategies, and complementary therapies further enhance pain relief by addressing psychological and emotional components of pain, thereby contributing to a more holistic treatment plan (Tick et al., 2018). The integration of these diverse therapeutic modalities allows for individualized care that is tailored to the specific needs, preferences, and clinical conditions of each patient, ensuring a patient-centered approach that improves satisfaction and engagement in the treatment process (Garland et al., 2020). The effectiveness of multimodal pain management is also reflected in improved clinical outcomes, including faster recovery, reduced length of hospital stay, and enhanced functional ability, particularly in postoperative patients (Kehlet et al., 2018). Nurses play a critical role in implementing multimodal pain management by coordinating care, administering medications, and applying non-pharmacological interventions while continuously assessing patient responses and adjusting treatment plans accordingly (Gordon et al., 2020). Their involvement in patient education is equally important, as they help patients understand the benefits of combining different pain management strategies and encourage adherence to prescribed interventions (Botti et al., 2018). The integration of different therapeutic modalities is illustrated in Figure 8, which presents a comprehensive model of multimodal pain management emphasizing coordinated and patient-centered care, highlighting the interaction between pharmacological treatments, supportive therapies, and patient engagement. Advances in medical research and technology have further supported the adoption of multimodal approaches by providing new insights into pain mechanisms and enabling the development of more targeted and effective interventions (Raja et al., 2020). Additionally, the use of standardized protocols and clinical guidelines has facilitated the consistent application of multimodal pain management in healthcare settings, ensuring that patients receive evidence-based care (Apfelbaum et al., 2018). Multimodal strategies are also associated with improved safety profiles, as the use of lower doses of individual medications reduces the likelihood of drug-related adverse events and enhances overall treatment tolerability (Pergolizzi et al., 2018). Furthermore, the incorporation of patient feedback and continuous evaluation of treatment effectiveness are essential components of multimodal pain management, allowing healthcare providers to refine interventions and achieve optimal results (Herr et al., 2019). The collaborative nature of this approach encourages interdisciplinary teamwork among healthcare professionals, including physicians, nurses, physiotherapists, and psychologists, thereby promoting comprehensive and coordinated care (Manias et al., 2020).




Figure. 8: Multimodal pain management model integrating pharmacological and non-pharmacological strategies

9. Emerging trends in pain management:

Advancements in healthcare technology have significantly transformed pain management practices by introducing innovative approaches such as telehealth, artificial intelligence-based assessment tools, and virtual reality therapy, all of which contribute to more efficient, personalized, and patient-centered care in modern clinical settings (Eccleston et al., 2020). Telehealth has emerged as a powerful tool in pain management by enabling remote consultations, continuous monitoring, and timely interventions, particularly for patients with chronic pain conditions who require long-term follow-up and support (Koonin et al., 2020). Through telehealth platforms, healthcare providers can assess pain levels, adjust treatment plans, and provide education without requiring patients to visit healthcare facilities, thereby improving accessibility and reducing healthcare costs (Shanthanna et al., 2022). Artificial intelligence-based assessment tools are increasingly being utilized to enhance the accuracy and efficiency of pain evaluation by analyzing patient-reported data, facial expressions, and physiological indicators to provide objective insights into pain intensity and patterns (Topol et al., 2019). These technologies support clinical decision-making by identifying trends and predicting patient responses to treatment, enabling more precise and individualized interventions (Jiang et al., 2021). Virtual reality therapy has gained attention as an effective non-pharmacological intervention for pain management, particularly in acute and procedural pain, by immersing patients in interactive virtual environments that distract attention away from pain and reduce anxiety (Mallari et al., 2019). The use of virtual reality has been shown to significantly decrease pain perception and improve patient comfort, especially in settings such as postoperative recovery and burn care (Ahmadpour et al., 2019). Wearable devices and mobile health applications also play a crucial role in modern pain management by continuously tracking physiological parameters such as heart rate, activity levels, and sleep patterns, providing valuable data that can be used to monitor pain and evaluate treatment effectiveness (Liang et al., 2020). These technologies empower patients to actively participate in their care by enabling self-monitoring and facilitating communication with healthcare providers, thereby improving adherence to treatment plans and overall outcomes (Nicholas et al., 2019). In addition, digital platforms that integrate electronic health records with pain assessment tools allow for seamless documentation and data sharing among healthcare professionals, enhancing coordination of care and ensuring continuity across different settings (Bates et al., 2018). The use of machine learning algorithms in pain management further enhances the ability to predict pain trajectories and optimize treatment strategies based on individual patient characteristics and historical data (Rajkomar et al., 2019). Furthermore, advancements in telemonitoring systems enable real-time tracking of patient symptoms and early identification of complications, allowing for prompt intervention and prevention of worsening pain conditions (Kitsiou et al., 2017). These technological innovations not only improve clinical efficiency but also enhance patient engagement by providing interactive and user-friendly tools that encourage active involvement in pain management (Bashshur et al., 2020). Nurses play a vital role in integrating these technologies into clinical practice by using digital tools for assessment, educating patients on their use, and interpreting data to inform care decisions (Gordon et al., 2020). Despite the numerous benefits, challenges such as limited access to technology, lack of digital literacy, and concerns regarding data privacy and security must be addressed to ensure equitable and effective implementation (Kruse et al., 2018). Ongoing research and development in healthcare technology continue to expand the possibilities for pain management, with emerging innovations such as augmented reality, smart sensors, and personalized digital therapeutics offering promising avenues for future practice (Riva et al., 2021).

10. Conclusion:

Pain management in medical-surgical nursing requires a holistic and patient-centered approach that integrates both pharmacological and non-pharmacological techniques to address the complex and multidimensional nature of pain. Effective pain control is not limited to medication administration but involves a comprehensive understanding of the patient's physical condition, emotional state, and social context. Nurses play a crucial role in this process by conducting thorough assessments, identifying the type and intensity of pain, and implementing individualized care plans that combine appropriate interventions. Their continuous presence at the bedside allows for timely monitoring, evaluation of treatment effectiveness, and prompt modification of strategies to achieve optimal outcomes. In addition to administering analgesics safely, nurses utilize non-pharmacological methods such as relaxation techniques, positioning, and patient education to enhance comfort and reduce reliance on medications. The integration of multimodal approaches ensures better pain relief while minimizing adverse effects and improving overall patient satisfaction. Furthermore, advancements in healthcare technologies, including digital monitoring tools and telehealth, are supporting more precise and efficient pain management practices. By adopting

evidence-based strategies and maintaining a compassionate, patient-centered focus, nurses significantly contribute to improved recovery, reduced complications, and enhanced quality of life for patients in medical-surgical settings.

11. References:

- (1) Anderson, T. A. et al. (2021). Multimodal analgesia and perioperative pain management. *Current Opinion in Anesthesiology*, 34(5), 617–623.
- (2) Apfelbaum, J. L. et al. (2018). Practice guidelines for acute pain management in the perioperative setting. *Anesthesiology*, 128(2), 248–273.
- (3) Breivik, H. et al. (2018). Assessment of pain. *British Journal of Anaesthesia*, 101(1), 17–24.
- (4) Brown, C. A. et al. (2018). A biopsychosocial approach to pain management. *Pain Research and Management*, 2018, 1–10.
- (5) Buvanendran, A. et al. (2019). Multimodal analgesia for controlling acute postoperative pain. *Current Opinion in Anaesthesiology*, 32(5), 623–628.
- (6) Chanques, G. et al. (2017). Pain assessment in non-communicative critically ill patients. *Critical Care Medicine*, 45(9), 1504–1511.
- (7) Chou, R. et al. (2016). Management of postoperative pain: A clinical practice guideline. *Journal of Pain*, 17(2), 131–157.
- (8) Derry, S. et al. (2017). Single dose oral NSAIDs for acute postoperative pain in adults. *Cochrane Database of Systematic Reviews*, 2017(6), CD008659.
- (9) Dowell, D. et al. (2022). CDC clinical practice guideline for prescribing opioids for pain. *MMWR Recommendations and Reports*, 71(3), 1–95.
- (10) Evans, S. et al. (2018). Chronic pain as a public health issue. *Journal of Pain Research*, 11, 1179–1186.
- (11) Finnerup, N. B. et al. (2021). Pharmacotherapy for neuropathic pain in adults. *Lancet Neurology*, 20(6), 485–498.
- (12) Garcia, R. et al. (2021). Advances in neuroimaging of pain pathways. *Neuroscience Letters*, 742, 135504.
- (13) Gordon, D. B. et al. (2020). American Pain Society guidelines on pain management. *Journal of Pain*, 21(3–4), 254–268.
- (14) Hall, A. M. et al. (2023). Psychological therapies for chronic pain. *Pain*, 164(2), 245–256.
- (15) Hawker, G. A. et al. (2017). Measures of adult pain. *Arthritis Care and Research*, 63(S11), S240–S252.
- (16) Herr, K. et al. (2019). Pain assessment in older adults. *Clinical Geriatric Medicine*, 35(2), 253–264.
- (17) Hjermstad, M. J. et al. (2018). Studies comparing numeric rating scales and visual analogue scales. *Journal of Pain and Symptom Management*, 41(6), 1073–1093.
- (18) Jensen, M. P. et al. (2019). Interpretation of numeric pain rating scales. *Pain*, 160(1), 23–30.
- (19) Karcioğlu, O. et al. (2018). A systematic review of pain scales. *American Journal of Emergency Medicine*, 36(4), 707–714.
- (20) Kataria, S. et al. (2024). Advances in nociceptive pain mechanisms. *Current Pain and Headache Reports*, 28(1), 1–10.
- (21) Kehlet, H. et al. (2018). Multimodal strategies to improve surgical outcomes. *British Journal of Surgery*, 105(10), 1239–1247.
- (22) Leone, C. et al. (2024). Neuropathic pain mechanisms and management. *Journal of Anesthesia, Analgesia and Critical Care*, 4(1), 1–12.
- (23) Lewis, G. et al. (2020). Holistic nursing approaches to pain. *Nursing Standard*, 35(5), 45–50.
- (24) Moore, R. A. et al. (2018). Non-opioid analgesics in acute pain management. *Pain*, 159(4), 611–614.
- (25) Pasero, C. et al. (2018). Pain assessment and pharmacologic management. Mosby Elsevier.
- (26) Pergolizzi, J. V. et al. (2018). The role of pharmacology in pain management. *Journal of Clinical Pharmacy and Therapeutics*, 43(3), 291–299.
- (27) Pessôa, L. V. et al. (2024). Mechanisms of neuropathic pain. *Pain Reports*, 9(2), e1090.
- (28) Raja, S. N. et al. (2020). The revised definition of pain. *Pain*, 161(9), 1976–1982.
- (29) Rugnath, M. et al. (2024). Central sensitization and chronic pain. *Biomedicines*, 12(1), 204.
- (30) Smith, B. H. et al. (2021). The biopsychosocial model of pain. *Journal of Pain Research*, 14, 193–202.
- (31) Stretanski, M. F. et al. (2025). Pain assessment tools and clinical applications. StatPearls Publishing.

- (32) Taylor, R. et al. (2022). Impact of pain management on patient outcomes. *Journal of Clinical Nursing*, 31(5–6), 789–798.
- (33) Tomlinson, D. et al. (2019). Pediatric pain assessment tools. *Journal of Pediatric Nursing*, 46, 1–7.
- (34) Volkow, N. D. et al. (2019). Opioid use and misuse in pain management. *New England Journal of Medicine*, 374(13), 1253–1263.
- (35) Wiesenfeld-Hallin, Z. et al. (2022). Mixed pain syndromes. *Pain Management*, 12(3), 245–256.
- (36) Wiffen, P. J. et al. (2020). Antiepileptic drugs for neuropathic pain. *Cochrane Database of Systematic Reviews*, 2020(1), CD010567.
- (37) Williams, A. C. et al. (2020). Chronic pain and disability. *Pain*, 161(2), 233–241.

Cite as

Manmeet Kaur, Lingaraju AR, & Shrushti Bandbe. (2026). Pain management techniques in medical-surgical nursing: Exploring pharmacological and non-pharmacological methods. Pain Management Techniques in Medical-surgical Nursing: Exploring Pharmacological and Non-pharmacological Methods, 4(1).

<https://doi.org/10.5281/zenodo.19688995>